

# Quantifying Some of the Major Sources of Uncertainty Associated with Estimates of Harp Seal Prey Consumption. Part II: Uncertainty in Consumption Estimates Associated with Population Size, Residency, Energy Requirement and Diet

**Peter A. Shelton, William G. Warren, and Garry B. Stenson**

Science Branch, Dept. of Fisheries and Oceans, Northwest Atlantic Fisheries Centre  
P. O. Box. 5667, St John's, Newfoundland, Canada A1C 5X1

and

**John W. Lawson**

Ocean Sciences Centre, Memorial University of Newfoundland  
St John's, Newfoundland, Canada A1C 5S7

## Abstract

Prey consumption by Northwest Atlantic harp seals, *Phoca groenlandica*, depends on population size, seasonal and spatial distribution, energy requirements, energy content of prey and diet composition. There is uncertainty in our knowledge of all these components. This carries through into uncertainty in any estimate of prey consumption. Available information ranges from sample estimates, sometimes with conventional measures of precision (standard errors), to guesses based on unquantified observation. An attempt is made here to quantify the effect of some of the major sources of uncertainty, particularly with respect to the amount of Atlantic cod (*Gadus morhua*) eaten in NAFO Division 2J and 3KL (off southern Labrador and northeast Newfoundland). The primary objective is to determine which components contribute most to the uncertainty, as a guide for research planning. However, a thorough quantification of uncertainty would also be useful in evaluating alternative management options for harp seals which have the objective of reducing possible impacts on prey. This work examines the effect on consumption estimates of uncertainty associated with the harp seal population size as well as the effect of additional sources of uncertainty attributable to residency of harp seals, energy requirements, species composition of the diet in the inshore and offshore, and the calorific value of prey.

*Key words:* consumption, Monte Carlo simulation, harp seal, uncertainty

## Introduction

The harp seal (*Phoca groenlandica*) population in the Northwest Atlantic was estimated to number 4.8 million in 1994 and to be increasing at about 5% per year (Shelton *et al.*, 1996). There is considerable interest in what impact this might have on fish populations in the region, particularly northern cod which is presently (since 1992) under a fishing moratorium because of low abundance. The estimated harp seal population size trajectory for the period 1981 to 1994 from Shelton *et al.* (1996) was used in Stenson *et al.* (1997) to calculate the consumption of Atlantic cod (*Gadus morhua*), capelin (*Mallotus villosus*) and Arctic cod

(*Boreogadus saida*). Fixed inputs were used for residency time, energy requirements and diet. From this analysis it was concluded that the harp seal population at present annually consumes 2.8 million tons of marine organisms from the Newfoundland region (NAFO Divisions 2J and 3KL) of which 1.7 million tons is Arctic cod, 0.6 million tons capelin and 0.09 million tons Atlantic cod. A preliminary sensitivity analysis was carried out in Stenson *et al.* (1997) by examining the effect of alternative plausible values for four key inputs. Changes in consumption values of up to 25% were obtained. Thus it was acknowledged that estimates of consumption are quite sensitive to changes in these inputs.

The purpose of the present work is to quantify the major sources of uncertainty in the calculation of consumption, primarily as a guide to where future research effort should be concentrated. However, a thorough quantification of uncertainty would also be useful in evaluating alternative management procedures for harp seals which have the objective of reducing possible impacts on prey. The robustness of alternative procedures to uncertainty, evaluated through simulation studies, should play a major role in selecting the best procedure. Although this study does not achieve the second objective, it attempts to lay the basis for further work in this direction.

A preliminary quantification of the uncertainty associated with the estimation of harp seal population size was carried out in Shelton *et al.* (1996). This is examined in much more detail in Warren *et al.* (1997). The present study examines further the uncertainty in the calculation of consumption contributed by the uncertainty in (i) population size, (ii) residency time in the study area, (iii) energy requirements, and (iv) diet, including species composition, energy content and assimilation efficiency. Conclusions are then drawn regarding where future research efforts should be concentrated in order to reduced uncertainty in the estimation of consumption. Consideration is also given to what further work needs to be carried out to arrive at reliable probability distributions of consumption by harp seals for use in decision making. Where the uncertainty is poorly known, the present analysis may be considered to be more akin to a sensitivity analysis. Reliable probability distributions of consumption will require greater information on the uncertainty with regard to the inputs.

## Methods

The uncertainty in the inputs was examined by Monte Carlo simulation. A single realization of consumption was generated by randomly selecting values for the inputs from distributions considered to describe the uncertainty in these inputs. This was repeated to give a set of realizations of consumption from which the effect of the factors being considered can be gauged. The analysis was conducted systematically, looking at each of the four main sources of uncertainty (population size, residency, energy requirement and diet) in turn, while holding all other inputs fixed at the values used in Stenson *et al.* (1997). Consumption was calculated for the period 1981 to 1994 (the period

for which diet data were available) and 50 randomly selected trajectories of consumption of cod plotted. Frequency distributions for 500 realizations of the 1994 consumption of cod, capelin and Arctic cod were plotted and basic univariate statistics tabled. Following the analysis of the contribution to the uncertainty by the four different sources individually, all modelled sources of uncertainty were included and 1 000 realizations were generated to get an impression of the possible overall uncertainty in the calculation of cod consumption. In order to examine the improvements that could be obtained with less certain inputs, each of the four sources of uncertainty were in turn treated as known exactly, while the remaining three sources were treated as uncertain and 500 realizations were generated. A more detailed analysis of uncertainty is possible by examining the individual contributors to uncertainty within each of the four sources and this could be considered in the future.

## Sources of Uncertainty

**Population size.** In Shelton *et al.* (1996) the uncertainty in the harp seal population trajectory was examined by randomly sampling pairs of parameters (survival rate and selectivity) from a bivariate normal distribution defined by the parameter estimates, their standard errors and the correlation between the estimates. It was acknowledged that this provided only a partial exploration of the uncertainty because it was assumed that pregnancy rates and catches were known exactly. It was therefore considered an underestimate of the uncertainty. Also, the viability of the asymptotic standard errors from only six years of survey data is questionable.

Warren *et al.* (1997) examined the uncertainty in the estimation of population size in much more detail. The uncertainty in both the estimates of pup production and the estimates of pregnancy rates were considered using an alternative non-asymptotic approach. It was found that the uncertainty in pregnancy rates had little additional effect when the uncertainty in population size was first accounted for. Results from the simulations to look at the uncertainty in pup production and the simulations in which both pup production and pregnancy rates are varied were combined to give 200 realizations of the population model parameter values. From these realizations Warren *et al.* (1997) developed a formulation from which pairs of parameter values can be randomly generated. First a realization of natural mortality,  $m$ , was generated

from a normal distribution with mean and standard deviation estimated from the 200 realizations:

$$m \sim N[0.106880, 0.007228]$$

Then a value of  $s$  was generated from a normal distribution with a mean  $\phi = 3.1219 - 30.3040m + 264.3880m^2$ :

$$s \sim N[\phi, 0.004073]$$

Note that in the notation used for a random variate from a normal distribution the first value in the square brackets is the mean and the second value is the variance.

As discussed in Warren *et al.* (1997) the problem with the non-asymptotic approach in which both the uncertainty in pregnancy rates and pup production are examined, is that the realizations of pregnancy rate need to be carried forward to the calculation of population size from the realization of the parameters of  $s$  and  $m$  obtained using those pregnancy rates. This is not possible in the parametric approach adopted here. It would require a full numerical simulation in which in each run a realization of pregnancy rate and a realization of pup production are generated from their respective probability distributions, the model is fitted and the realizations of the estimated parameters  $s$  and  $m$  used with the identical realization of pregnancy rate to generate a population trajectory. This approach was considered impractical for the present study but warrants future attention. Since Warren *et al.* (1997) found that uncertainty in pregnancy rates inflates the overall uncertainty in population size marginally, the parametric approach was applied here with constant pregnancy rates with the knowledge that the uncertainty in population size was underestimated by some amount.

**Residency time.** Harp seals summer in Arctic waters and winter off Newfoundland and Labrador and in the Gulf of St. Lawrence (Sergeant, 1965, 1991). The general migration pattern has been determined from surveys, catches, aerial observations and anecdotal sightings, however detailed knowledge is limited (Stenson *et al.* 1997). Uncertainty in the residency time was examined in a preliminary manner in Stenson *et al.* (1997) by calculating consumption based on a fixed residency of 212 days (south of the northern boundary of Div. 2J) and then comparing this with a one month increase in residency. The increase in residency increased consumption by 12%. In this analysis we look in

more detail at the uncertainty in the various factors that must be taken into account in determining the residency within Divisions 2J and 3KL and how this affects the calculation of consumption of Atlantic cod, capelin and Arctic cod.

Harp seals migrate south from the summer feeding grounds in the Arctic during the late autumn. Based on catches and sightings summarized by Sergeant (1965, 1991), Stenson *et al.* (1997) assumed that the average date seals entered the study area (south of the Divisions 2J/2H boundary) was 15 November and that they left on 15 June. However, the migration may be spread over a relatively long period (Sergeant, 1965) and the timing of the peak migration may vary greatly (Fisher 1955; Stenson, NWAFC, St. John's, Newfoundland, unpubl. data) with reports of seals within the study area from early October through July. Therefore, to quantify the uncertainty associated with this parameter, it was assumed that seals may enter the study area between 15 October and 1 December and leave between 1 June and 15 July. This gives a range of possible residency time within the study area of between 182 and 272 days, as compared to the fixed value of 212 days used in Stenson *et al.* (1997). A uniform distribution within this range was assumed. Once in the study area, a proportion of the population enters the Gulf of St. Lawrence after migrating through Divisions 2J and 3KL. Stenson *et al.* (1997) assumed a period of half a month for each of the southern and northward migrations (total 29 days), based upon the respective timing of fisheries along the mid-Labrador and northern Quebec coasts (Sergeant, 1991). However, movements of individual seals obtained using satellite telemetry indicate that harp seals may move quickly between areas and then remain in one location for a considerable time. Therefore, the timing of the migration through the Newfoundland area may vary greatly. To quantify the uncertainty associated with this assumption, the amount of time animals destined for the Gulf spend within the study area was assumed to be described by a uniform distribution of between 15 and 45 days. Stenson *et al.* (1997) assumed that the proportion of the population entering the Gulf was 0.25. This was based upon the assumption that approximately 1/3 of adults enter the Gulf to whelp while some immatures remain off Newfoundland (Sergeant, 1991). However, the proportion of total pup production which occurs in the Gulf can vary greatly among years. Comparing estimates obtained from comparable aerial surveys of both areas indicates that the proportion of total pup

production which occurred in the Gulf rose from approximately 0.19 in 1990 (Stenson *et al.*, 1993) to 0.34 in 1994 (Stenson *et al.*, 1997). Similarly, Winters (1978) estimated that the proportion of the total annual pup production which occurred in the Gulf from 1965–77 varied between 0.13 and 0.51. Therefore, a uniform distribution within the range of 0.2 to 0.4 was assumed. A proportion of the population remains in the Arctic throughout the year. A range of 0.15 to 0.25 was assumed compared with a fixed value of 0.2 used in Stenson *et al.* (1997)

### Energy requirement

Stenson *et al.* (1997) calculated individual energy requirements using an allometric relationship linked to mass-at-age based on Kleiber (1975). Corrections for the additional energy requirements associated with growth, activity and assimilation efficiency are incorporated. The energy requirements for individual harp seals is assumed to be constant throughout the year. The equation is:

$$GEI_i = GP_i \times (AF \times 70 \times BM_i^{0.75}) / (ME)$$

where  $GEI$  is the daily gross energy intake,  $i$  is the age group,  $GP$  is the growth premium (energy cost of growth),  $AF$  is the "activity factor",  $BM$  is the mean body mass for age group (kg), and  $ME$  is the proportion of energy available to the animal (assimilation efficiency).

Body mass for each age group was based on measurements obtained from seals collected during April (Chabot *et al.*, 1996). To account for uncertainty in body size we randomly resampled body mass-at-age values from a normal distribution defined by the mean and standard deviation of these sample data:

Age	Value
0	$N(25.449, 5.442)$
1	$N(45.846, 7.279)$
2	$N(56.041, 10.064)$
3	$N(64.755, 10.354)$
4	$N(74.863, 13.988)$
5	$N(82.278, 13.648)$
6	$N(85.384, 12.703)$
7	$N(92.783, 12.563)$
8	$N(93.487, 13.971)$
9	$N(96.504, 13.958)$
10	$N(101.763, 13.128)$
11	$N(101.763, 13.128)$
12+	$N(101.763, 13.128)$

Lavigne *et al.* (1986) reviewed literature on metabolic rates of seals and suggested that growing phocids had basal metabolic rates twice that of older animals. The increased energy required for growth was applied to the metabolic calculations for younger seals in decreasing increments from  $GP = 2.25$  for one-year olds to 1.25 for 5 year olds. These values were used in Stenson *et al.* (1997). Alternative values are given in Olesiuk (1993). In the present analysis, each realization randomly selects between the values given by Lavigne *et al.* (1986) and those given by Olesiuk (1993) with equal probability. Further information on this input may allow in the future a more comprehensive examination of the uncertainty contributed by this source.

Studies of the energy requirement of captive and wild seals indicate that estimates of the average daily energy requirements vary between 1.7 and 3 times the basal metabolic rate estimated using body mass (Castellini *et al.*, 1992; Innes *et al.*, 1987; Worthy, 1987a; 1987b; 1990). Since most published values cluster near a value of 2, an activity factor ( $AF$ ) of 2 was chosen in Stenson *et al.* (1997) to approximate the energy requirements of activity of free-ranging harp seals. To account for the uncertainty, activity factors were randomly sampled from a triangular distribution. The distribution extended from 1.7 to 3 and had a peak at 2. The area of the triangle was made to sum to 1 by setting the height to 1.5385. A rejection method was used to randomly sample from this distribution. Two random variates,  $X \sim U[1.7, 3]$  and  $Y \sim [0, 1.5385]$  were generated in each realization until the coordinates were within the area of the triangle. The  $X$  value for realizations within the triangle was taken to be a realization of the activity factor.

### Diet

Considerable seasonal, geographic, and annual variability exists in the diet of harp seals (Lawson *et al.*, 1995; Stenson *et al.* 1997; Lawson and Stenson, 1995). Stenson *et al.*, (1997) presented information on the diet of harp seals in Newfoundland separated into winter and summer periods for offshore diets (1991–94 combined) and 6 years for which reconstructed stomach contents of nearshore harp seals were available (1982, 1986, 1990–93). To express the uncertainty in the proportion of prey in the diet, they estimated consumption by harp seals in Newfoundland based on the annual diets as well as consumption estimated using the average of the 14 annual diet averages. The uncertainty associated with using the

overall average diet was illustrated by calculating the 95% confidence intervals around the estimates of consumption by randomly resampling the 14 diets with replacement 1 000 times.

In the present analysis the stomach sample data were treated in two different ways. In the first treatment, Diet A, composition by weight samples for the years 1990 to 1993 were separated into summer and winter groups for the inshore and offshore (i.e. a total of 4 groups). Samples within each of these four groups were found to be relatively homogeneous. A realization of the diet composition from Diet A was obtained by randomly selecting diet samples with replacement from the individual diet samples contained in that group until the number of diets selected equals the sample size of that group. The average diet composition in each group was then obtained by dividing the group total by the sample size for that group. The contribution of each diet to the overall estimated of annual consumption was weighted according to the weightings representing the relative amount of energy obtained from each of the four sets taken from Stenson *et al.* (1997). The relative weightings used for winter:summer were 0.6241:0.3759 and for inshore:offshore were 0.45:0.55.

Alternative analyses were carried out using Diet B for comparison with the results of Stenson *et al.* (1997). In this treatment diet data from the inshore for 1982, 1986 and 1990–93, were separated into summer and winter groups for each year (see Stenson *et al.* 1997). Offshore data from all years combined were also separated into summer and winter groups. This gives a total of 14 groups overall. A realization from Diet B was obtained by randomly selecting individual diet samples with replacement from within each group until the number selected equaled the sample size for that group. The within group average diet from the random resampling was then obtained by summing within group and dividing through by the sample size. Finally, the overall average diet composition was obtained, applying the same weightings as in the treatment using Diet A.

In the analysis carried out introducing each source of uncertainty alone, Diet B was used (i.e. fixed as per Stenson *et al.* (1997)). However, when diet itself was varied, then random realizations from Diet A are generated. When all source of uncertainty were introduced together and then each source removed one at a time, Diet A was used. A final analysis including all sources of uncertainty introduced

except uncertainty in diet was carried out but with diet set according to Diet B.

The proportion of energy contained in the food which is available to the harp seals (assimilation efficiency) has not been measured directly for most of the major prey items. Published values range from 72.2% for shrimp (Keiver *et al.*, 1984) to 94% for capelin (Mårtensson *et al.*, 1994). Following Stenson *et al.* (1997) the mean assimilation efficiency were calculated here for each of the four realized sets of diet data using fixed prey-specific values. Energy density of prey were based on the average of published values for the major species, where available (e.g. Anon., 1969, Croxall and Prince, 1982, Hislop *et al.*, 1991, Holdway and Beamish, 1984, Hop, 1994, Hopkins *et al.*, MS 1989; Krzynowek and Murphy, 1987; Nordøy and Blix, 1988; Steimle Jr. and Terranova, 1985) and analyses performed at Northwest Atlantic Fisheries Centre, Newfoundland (Lawson, unpubl. data). Following Stenson *et al.* (1997) the mean energy content of the prey was calculated based on the diet composition.

It should be noted that this treatment of the diet data in the present analysis is different to that of Stenson *et al.* (1997). In particular, the 1982 and 1986 diet data, which are significantly different to the more recent diet data (Warren, unpubl. analysis), are not used in the present analysis. Further, the diet data from 1990 onwards, within each of the four sets, are treated as if they are all equally likely realizations of the 1994 diet (i.e. there have been no systematic changes over the period). In the calculation of a cod consumption trajectory, the assumption is made that the realization of diet that is randomly selected from the diet samples for the period 1990–93 applies over the period 1981 to 1994. This assumption is unlikely to be valid.

## Results and Discussion

The 50 realizations of cod consumption over the period 1981 to 1994 shows the overall increase caused by the increase in the seal population (Fig. 1). The relative contribution to the uncertainty from the four sources as illustrated in these plots shows that uncertainty in population size has the least effect and uncertainty in diet the greatest effect. The contribution by uncertainty in residency and energy requirements are similar. Note that the uncertainty in diet examined here follows the approach described above (resampling from four relative homogeneous sets of the data for the post




Fig. 1. Plots of 50 realizations of cod consumption trajectories for the period 1981 to 1994 taking into account uncertainty in population size, residency, energy requirements and diet.

1990 period) whereas the diet is fixed at the values given in Stenson *et al.* (1997) in the other three analyses.

The uncertainty in consumption of cod, capelin and Arctic cod contributed by each of the four sources of uncertainty alone are illustrated in the

frequency distributions for 500 realizations of consumption (Figs. 2–4). Descriptive statistics for the distribution of cod consumption are given in Table 1. With respect to cod consumption (Fig. 2, Table 1), uncertainty in diet made the greatest contribution to uncertainty in cod consumption ( $CV = 28\%$ ) and uncertainty in population size made the




Fig. 2. Frequency distributions of 500 realizations of cod consumption taking into account uncertainty in population size, residency, energy requirements and diet.




Fig. 3. Frequency distributions of 500 realizations of capelin consumption taking into account uncertainty in population size, residency, energy requirements and diet.

smallest contribution ( $CV = 4\%$ ). Uncertainty in energy requirement and residency made equal contributions ( $CV = 12\%$ ). It is of interest that the fifth percentile for analysis in which uncertainty in diet is accounted for is not any lower than in the other analyses, however the ninetieth percentile is much higher, giving a large 90% probability range. Recall that the distribution in which diet is varied does

not centre around the 88 000 tons value given in Stenson *et al.* (1997) because of the different treatment of the diet data.

With respect to capelin consumption (Fig. 3), uncertainty in population has a relatively small effect compared to uncertainty in the other sources. The spread in the distribution caused by uncertainty




Fig. 4. Frequency distributions of 500 realizations of Arctic cod consumption taking into account uncertainty in population size, residency, energy requirements and diet.

in residency and energy requirements are similar, whereas the spread resulting from uncertainty in diet is only slightly larger.

Uncertainty in Arctic cod consumption (Fig. 4) is relatively small when uncertainty in harp seal population size is taken into account. Uncertainty in both residency and energy requirements make

relatively large contributions whereas the contribution to uncertainty by diet is approximately the same as that of population size.

When all four sources of uncertainty are introduced simultaneously, there is a wide range in the amount of cod, capelin and Arctic cod consumed, as might be expected (Fig. 2–5, Table 1). In the 50

TABLE 1. Descriptive statistics for the distributions of 500 realizations of cod consumption including each source of uncertainty separately, all sources together, and then removing each source in turn. Note that "Diet-A" refers to the diet grouped into four sets representing inshore, offshore, summer and winter, and that "Diet-B" is the diet used in Stenson *et al.* (1997). The analysis introducing each source of uncertainty separately uses "Diet-B" except for when diet itself is varied, when "Diet-A" is used. When the sources of uncertainty are removed one at a time, 500 realizations are carried out and the results described for both diet treatments, to allow comparison between treatments and with Stenson *et al.* (1997). Diet A comprises data from the nearshore for 1990 to 1993 for both summer and winter and data combined across years for the offshore in both summer and winter, a total of 12 different categories.

Source	Mean	Median	Std Dev	CV	Min	Max	P5	P95	90% P range
<b>Source added</b>									
Population size	90137.42	90877.28	3662.75	4.06352	72440.62	121312.90	84032.11	93329.41	9297.30
Residency time	89348.25	89857.01	10544.04	11.8011	66250.81	116787.71	72801.78	106830.05	34028.27
Energy requirement	88169.91	86916.14	10533.78	11.9471	62452.92	118916.16	72976.57	106943.35	33966.78
Diet composition – A	128665.79	125876.11	35559.71	27.6373	49080.18	271084.94	72658.90	194145.19	121486.29
-----									
<b>All sources together</b>	140807.54	135128.17	49554.66	35.1313	49554.66	416255.59	72544.34	232100.10	159555.76
<b>Source removed</b>									
Population size	132334.32	125723.74	45965.76	34.7346	42643.16	284215.86	70475.86	225670.79	155194.93
Residency time	138386.73	134900.79	43994.29	31.7908	47679.57	303639.29	78898.51	216429.85	137531.34
Energy requirement	143947.34	140608.56	43419.02	30.1631	46892.69	330059.46	80260.37	219815.15	139554.78
Diet composition – A	130693.61	128340.73	23955.95	18.3299	79386.40	237204.76	95898.80	172637.94	76739.14
Diet composition – B	91818.32	90499.57	16144.72	17.5833	56766.03	149070.29	67878.98	120207.11	52328.13

realizations of cod consumption over the period 1981 to 1994, cod consumption ranged from about 30 000 tons to 150 000 tons in 1980 and about 50 000 tons to 300 000 tons in 1994. The distributions from the 1 000 realizations of consumption for cod, capelin and Arctic cod are all slightly skewed to the right, i.e. there is a small probability of consumption being quite a bit higher than the mean, but less probability that consumption is much smaller than the mean. The CV in cod consumption is 35% with fifth and ninetieth percentiles of 73 000 tons and 232 000 tons, respectively. Uncertainty in capelin consumption ranges from 360 000 tons to 1 500 000 tons and Arctic cod consumption from 560 000 tons to 1 700 000 tons.

The reduction in overall uncertainty in cod consumption obtained by, in turn, treating each of the four individual sources of uncertainty as known exactly is summarized in Table 1. As anticipated, knowing harp seal population size exactly gives the smallest improvement in the CV – from 35.1% to 34.7%. Improvements obtained by knowing either the residency time of harp seals in the study area or their energy requirements exactly are similar (CV

reduced to 31.7% and 30.2%, respectively). Exact information on diet reduces the CV on the uncertainty to 18.3%. A similar CV would pertain if the diet as used in Stenson *et al.* (1997) was used. Note that there is little improvement in the minimum estimate by removing uncertainty in any of the sources other than diet.

These results suggest that the largest improvements in the precision of estimates of harp seal consumption of cod will be obtained by improved knowledge of diet composition. This conclusion is based on the assumption that within each of the four relatively homogeneous diet sets for the period 1990 onwards, the samples vary according to random sampling error only and there is no systematic variability that can be accounted for (e.g. a time trend in diet in the period 1990 onwards).

The conclusion made in the case of cod consumption with respect to knowing diet exactly does not appear to pertain to capelin and Arctic cod consumption. For capelin, uncertainty in residency, energy requirements and diet contribute approximately equally, whereas in the case of Arctic cod,




Fig. 5. Plot of 50 realizations of cod consumption trajectories over the period 1981 to 1994 together with frequency distributions of cod, apelin and Arctic cod consumption from 1 000 realizations taking into account all four sources of uncertainty simultaneously.

uncertainty in residency and energy requirement contribute the most.

While better information for all four groups of inputs into the estimation of harp seal consumption

would be very valuable, the present study suggests that improvements in the diet data will yield the most benefit with respect to the improvements in the estimate of cod consumed by harp seals. Further work could be carried out at a finer scale by

examining the contribution of uncertainty by the various factors within each of the four sources. The present analysis only looks at the improvement obtained by going from a situation in which the input is uncertain to one in which it is known exactly. More detailed analysis could look at the effect of certain percentage reductions in the uncertainty of selected inputs and the financial cost of making such reductions. This would allow research planning of a more detailed kind than simply the suggestion that "we need better diet data".

It is of interest that the uncertainty in population size is the smallest contributor to uncertainty in consumption. It is also one of the few inputs for which a formal estimate of the uncertainty is available. Although intensification of the pup survey program through more frequent surveys may not lead to a large improvement in the uncertainty regarding harp seal consumption of prey species, the more time-consuming simulation approach described above should be explored to examine the potential improvement that may result from better pregnancy data. Although Warren *et al.*, (1997) suggests that the contribution by uncertainty in pregnancy rate is small, there are difficulties in interpreting the current data set because of the relative paucity of samples in some years.

Considerably more work remains to be done before a thorough quantification of the uncertainty regarding cod consumption can be arrived at for use in decision making. Nevertheless, in the interim, decision makers should be cognizant of the fact the CV in the current estimate of cod consumption is of the order of 35%. Over-reliance on a point estimate should be avoided when possible.

### Acknowledgements

This analysis is based on the estimation model constructed by Noel Cadigan and would not have been possible without his insights. Nicholas Barrowman is thanked for telling one of us about the rejection method for generating random values from a triangular probability distribution.

### References

- ANON. 1969. United States – Canadian tables of feed composition. Subcommittee on Feed Composition, *National Academy of Sciences*. 1684.
- CASTELLINI, M. A., G. L. KOOYMAN, and P. J. PONGANIS. 1992. Metabolic rates of freely diving Weddell seals: Correlations with oxygen stores, swim velocity and diving duration. *J. Exp. Biol.*, **165**: 181–194.
- CHABOT, D., G. B. STENSON, and N. G. CADIGAN. 1996. Short- and long-term fluctuations in the size and condition of harp seal (*Phoca groenlandica*) in the northwest Atlantic. *NAFO Sci. Coun. Studies*, **26**: 15–32.
- CROXALL, J. P., and P. A. PRINCE. 1982. Calorific content of squid (Mollusca: Cephalopoda). *British Antarctic Survey Bulletin*, **55**: 27–31.
- FISHER, H. D. 1955. Utilization of Atlantic harp seal populations. *Trans. North Amer. Wildl. Conf.*, **20**: 507–518.
- HISLOP, J. R. G., M. P. HARRIS, and J. G. M. SMITH. 1991. Variation in the calorific value and total energy content of the lesser sandeel (*Ammodytes marinus*) and other fish preyed on by seabirds. *J. Zool.*, London, **224**: 501–517.
- HOLDWAY, D. A., and F. W. H. BEAMISH. 1984. Specific growth rate and proximate composition of Atlantic cod (*Gadus morhua* L.). *J. Exp. Mar. Biol.*, **81**: 147–170.
- HOP, H. 1994. Bioenergetics and trophic relationships of Arctic cod (*Boreogadus saida*) in the Canadian High Arctic. Ph.D. Thesis, University of Alberta, Edmonton.
- HOPKINS, C. C. E., E. M. NILSSEN, A. HERMANNSEN, B. VAAJA, and L. DALSBØE. MS 1989. Body composition and energetic predictors of age/season in the prawn *Pandalus borealis*. *ICES C. M. Doc.*, No. K:10.
- INNES, S., D. M. LAVIGNE, W. M. EARLE, and K. M. KOVACS. 1987. Feeding rates of seals and whales. *J. Anim. Ecol.*, **56**: 115–130.
- KEIVER, K. M., K. RONALD, and F. W. H. BEAMISH. 1984. Metabolizable energy requirements for maintenance and faecal and urinary losses of juvenile harp seals (*Phoca groenlandica*). *Can. J. Zool.*, **62**: 679–776.
- KLEIBER, M. 1975. The fire of life: An introduction to animal energetics. New York: Robert E. Krieger Publishing Co.
- KRZYNOWEK, J., and J. MURPHY. 1987. Proximate composition, energy, fatty acid, sodium, and cholesterol content of finfish, shellfish, and their products. *NOAA Tech. Rep.*, NMFS 55.
- LAVIGNE, D. M., S. INNES, G. A. WORTHY, and K. M. KOVACS. 1986. Metabolic rate-body size relations in marine mammals. *J. Theor. Biol.*, **122**: 123–124.
- LAWSON, J. W., and G. B. STENSON. 1995. Historic variation in the diet of harp seals (*Phoca groenlandica*) in the northwest Atlantic. In: Whales, seals fish and man. A.S. Blix, L. Walløe and Ø. Ultang (eds.) Elsevier Science, 261–269 p.
- LAWSON, J. W., G. B. STENSON, and D. G. MCKINNON. 1995. Diet of harp seals (*Phoca groenlandica*) in nearshore waters of the northwest Atlantic during 1990–1993. *Can. J. Zool.*, **73**: 1805–1818.
- MÅRTENSSON, P. E., E. S. NORDØY, and A. S. BLIX. 1994. Digestibility of crustaceans and capelin in

- harp seals (*Phoca groenlandica*). *Mar. Mam. Sci.*, **10**: 325–331.
- NORDØY, E.S., and A.S. Blix. 1988. Sources of error in estimating food requirements of seals. *Polar Record*, **24**: 62–64.
- OLESIUK, P. F. 1993. Annual prey consumption by harbour seals (*Phoca vitulina*) in the Strait of Georgia, British Columbia. *Fish. Bull.* (Dublin), **91**: 491–515.
- SERGEANT, D. E. 1965. Migration of harp seals *Pagophilus groenlandicus* (Erxleben) in the Northwest Atlantic. *J. Fish. Res. Board Can.*, **22**: 433–463.
- SERGEANT, D. E. 1991. Harp seals, man and ice. *Can. Spec. Pub. Fish. Aquat. Sci.*, **114**, 153p.
- SHELTON, P. A., G. B. STENSON, B. SJARE, and W. G. WARREN. 1996. Model estimates of harp seal numbers-at-age for the Northwest Atlantic. *NAFO Sci. Coun. Studies*, **26**: 1–14.
- STEIMLE Jr., F. W., and R. J. TERRANOVA. 1985. Energy equivalents of marine organisms from the continental shelf of the temperate Northwest Atlantic. *J. Northwest Atl. Fish. Sci.*, **6**: 117–124.
- STENSON, G. B., R. A. MYERS, M. O. HAMMILL, I-H. NI, W. G. WARREN, and M. C. S. KINGSLEY. 1993. Pup production of harp seals, *Phoca groenlandica*, in the Northwest Atlantic. *NAFO Sci. Coun. Studies*, **26**: 47–61.
- STENSON, G. B., M. O. HAMMILL, M. C. S. KINGSLEY, B. SJARE, W. G. WARREN, and R. A. MYERS. MS 1996. Pup production of Northwest Atlantic harp seal, *Phoca groenlandica*. *DFO Atl. Fish. Res. Doc.*, No. 20.
- STENSON, G. B., M. O. HAMMILL, and J. W. LAWSON. 1997. Predation by harp seals in Atlantic Canada: preliminary consumption estimates for Arctic cod, capelin and Atlantic cod. *J. Northwest Atl. Fish. Sci.*, **22**: 137–154.
- WARREN, W. G., P. A. SHELTON, and G. B. STENSON. 1997. Quantifying some of the major sources of uncertainty associated with estimates of harp seal prey consumption. Part I: Uncertainty in estimates of harp seal population size. *J. Northwest Atl. Fish. Sci.*, **22**: 289–302.
- WINTERS, G. H. 1978. Production, mortality, and sustainable yield of northwest Atlantic harp seals (*Pagophilus groenlandica*). *J. Fish. Res. Board Can.*, **35**: 1249–1261.
- WORTHY, G. A. J. 1987a. Ecological energetics of harp and gray seals: Implications from a simulation model. *In: Marine Mammal Energetics*, A. C. Huntley, D. P. Costa, G. A. J. Worthy and M. A. Castellini (eds.). Allen Press, Lawrence, Kansas, p. 227–246.
- WORTHY, G. A. J. 1987b. Metabolism and growth of young harp and grey seals. *Can. J. Zool.*, **65**: 1377–1382.
- WORTHY, G. A. J. 1990. Nutritional energetics in marine mammals: Addendums. *In: CRC handbook of marine mammal medicine: Health, disease, and rehabilitation*, L.A. Dierauf (ed.), CRC Press, Boca Raton, Florida, p. 489–520.
-