

Biodiversity of Bear Seamount, New England Seamount Chain: Results of Exploratory Trawling

J. A. Moore

Florida Atlantic University, Honors College, Jupiter, FL 33458, USA

M. Vecchione, B. B. Collette and R. Gibbons

National Marine Fisheries Service, National Systematics Laboratory,
National Museum of Natural History, Smithsonian Institution, Washington, DC 20560

K. E. Hartel

Museum of Comparative Zoology, Harvard University, Cambridge, MA 02138, USA

J. K. Galbraith

National Marine Fisheries Service, Northeast Fisheries Science Center, Woods Hole, MA 02453, USA

M. Turnipseed, M. Southworth, and E. Watkins

Virginia Institute of Marine Sciences, Gloucester Point, VA 23062, USA

Abstract

Bear Seamount (39°55'N 67°30'W) is an extinct undersea volcano located inside the U.S. Exclusive Economic Zone south of Georges Bank. The fauna associated with the seamount was little known until twenty trawl stations were made 2–7 December 2000, by the NOAA ship R/V *Delaware II*. The objective of the survey was to begin to document the biodiversity on and over the seamount, particularly of fishes, cephalopods, and crustaceans. Representatives of most species were preserved as vouchers and for subsequent definitive identification. This report presents a description of the biodiversity. A total of at least 274 species were collected. Preliminary identifications indicate the capture of 115 fish species. Among these were a number of new fish records for the area or rare species, including *Acromycter pertubator* (Congridae), *Diastobranchius capensis* (Synaphobranchidae), *Alepocephalus bairdii* (Alepocephalidae), *Mirognathus normani* (Alepocephalidae), *Bathygadus favosus* (Bathygadidae), *Nezumia longebarbata* (Macrouridae), *Gaidropsarus argentatus* (Phycidae), *Dibranchius tremendus* (Ogcocephalidae) and *Kali indica* (Chiasmodontidae). Only two fish species of potential commercial importance were encountered: *Coryphaenoides rupestris* and *Macrourus berglax*. Cephalopods comprised 26 species from 15 families, including one new distributional record and several rarely-collected species. The crustacean fauna was diverse with at least 46 species. Totals for other invertebrate species are pending laboratory identification, but number at least 87 species from 10 Phyla. This includes a number of new distributional records and a new species of gorgonian.

Key words: biodiversity, deep-sea, fauna, fishes, invertebrates, North Atlantic, seamount

Introduction

Seamounts represent biological islands in the deep sea and often feature characteristic faunas that are quite different from those found in the surrounding soft sediment and abyssal habitats. Large depth ranges, hard substrates, steep gradients, complex topography, impinging currents, topographically induced upwellings, clear oceanic water, and geographic isolation all combine to make seamounts specialized habitats for deep-sea organisms (Rogers, 1994). The benthic fauna on seamounts include species that cre-

ate structural habitat diversity and likely create essential habitats for some deep-sea fishes and crustaceans.

The New England Seamounts (NES) make up the longest seamount chain in the North Atlantic, encompassing more than 30 major volcanic peaks extending southeast from Georges Bank for about 1 200 km to the eastern end of the Bermuda Rise, ending abruptly with Nashville Seamount to the East North East of Bermuda (Fig. 1). The major peaks of the NES rise as much as 4 000 m above the Sohm Abyssal Plain.

With the increase of deep-sea fisheries, interest in commercially fishing seamounts has increased (Koslow, 1997). Yet seamounts hold concentrations of sessile suspension-feeding invertebrates, such as scleractinians, gorgonians and sponges, that add structural complexity, and there are concerns about the impact of fishing activities on these invertebrates (Dayton *et al.*, 1995; Auster *et al.*, 1996; Probert *et al.*, 1997; Koslow and Gowlett-Holmes, MS 1998; Moore and Mace, 1999; Koslow *et al.*, 2000). A recent haul of Darwin's roughy (*Gephyroberyx darwini*, also known as big roughy) from a canyon on the south side of Georges Bank generated renewed interest by some U.S. fishers in exploring the deepwater fishery resources of New England. One fishing company put forward a specific proposal to explore the fishery resources on Bear Seamount. This coincided with planning by the National Marine Fisheries Service (NMFS) to conduct an exploratory survey to Bear Seamount in late-2000. One objective of this survey was to begin to document the biodiversity on and over the seamount, particularly of fishes, cephalopods, and crustaceans before large-scale commercial fishing begins. Knowledge of the fauna associated with the seamounts is considered crucial in assessing the impacts of any subsequent commercial fishing on these seamounts and a lack of this kind of information has hindered assessments of fishing impacts in other areas (Koslow and Gowlett-Holmes, MS 1998).

Bear Seamount (39°55'N 67°30'W) is an extinct undersea volcano located inside the U.S. Exclusive Economic Zone south of Lydonia Canyon on Georges Bank. Bear Seamount rises from the continental slope at depths of 2 000–3 000 m to a generally flat summit at a depth of 1 100 m below the sea surface. Although the upper surface of Bear Seamount is draped in many places with thick sediment, there are also outcrops of basaltic volcanic rock and scattered glacial erratics varying in size from small pebbles to large boulders. These glacial erratics most likely fell to the seamount after dropping from icebergs calving off the glaciers that extended to Georges Bank during the Pleistocene.

Bear Seamount was first identified, named and sonar mapped during a series of surveys by Woods Hole Oceanographic Institution (WHOI) vessels in 1954 (Zeigler, MS 1955). Subsequent investigations of Bear Seamount include more sonar mapping, one chain-bag dredge haul and one camera lowering in 1962 (R/V *Atlantis* cruise A-281; Pratt and Thompson,

MS 1962), two *Alvin* dives in 1968 (*Alvin* dive 286 and 287), and a commercial exploratory trawl by the F/V *Matthew Melissa* out of Stonington, Connecticut, in 1997. A list of taxa identified from these previous studies is given in Table 1.

Prior to our first cruise in 2000, several invertebrates and a few fishes were identified from archived photos from *Atlantis* cruise A-281 and *Alvin* dives 286 and 287. The archived photos are available at the Document Library of Woods Hole Oceanographic Institution. In addition, several fish specimens were acquired from a commercial trawler that conducted a few deepwater exploratory tows on the top of Bear Seamount in 1997. These specimens are deposited at the Ichthyology Collection of the Peabody Museum of Natural History, Yale University.

Materials and Methods

Twenty hauls were made on or over Bear Seamount using double-warp gear towed from the NOAA vessel R/V *Delaware II* from 2–7 December 2000 (cruise DE00-11). Of these hauls, 11 were fully pelagic tows and 9 actually touched bottom, if only briefly in some instances. Each tow was for one hour once the net was set at the selected depth, however, bottom tows were not necessarily in contact with the bottom that entire time. Tow depths were between 900–2 500 m. Six hauls were made with an IYGPT midwater trawl, five hauls with a NMFS standard shrimp trawl, and nine tows with a Yankee 36 otter trawl. The differing biases and catchabilities of these three kinds of nets, plus the unknown amount of time spent trawling the bottom, make explicit calculations of diversity indices suspect. Catches were sorted and given preliminary identifications on shipboard. Tissue samples of many fish and cephalopod taxa were taken for DNA analysis (samples of fishes are deposited at the Museum of Natural History, University of Kansas). Representatives of most species were preserved as vouchers. The voucher specimens are deposited in the Museum of Comparative Zoology at Harvard University (most of the fish specimens), Peabody Museum of Natural History at Yale University (some invertebrates), and the National Museum of Natural History, Smithsonian Institution (some fishes and most of the invertebrates). Many specimens were sent to taxonomic experts for subsequent definitive identification. A list of those experts is given in the acknowledgments.

TABLE 1. Taxa seen or sampled from Bear Seamount prior to 2000.

PROTISTA	
Globose syringaminid? xenophyophore	
PORIFERA	
Unidentified sponges (>5 spp.)	<i>Neomorphaster</i> sp. small 5-arm sea star <i>Ophiomusium lymani</i> brittle star Unidentified light brittle stars <i>Echinus affinis</i> urchin Echinothuriid pancake urchin Unidentified 5-arm bathyrcrinid crinoid <i>Peniagone</i> sp. sea cucumber
CNIDARIA	
Cerianthid anemone Unidentified large light-colored anemone <i>Desmophyllum cristagalli</i> scleractinian coral <i>Lepidisis</i> sp. unbranched spiral whip coral <i>Primnoa</i> sp. gorgonian coral Pinnate antipatherian corals <i>Anthophilum</i> sp. sea pen	UROCHORDATA
	Large transparent unstalked tunicate
SIPUNCULIDA	
Unidentified sipunculid worm	PISCES
ANNELIDA	
<i>Hyalinoecia</i> sp. polychaete worm	<i>Hydrolagus pallidus</i> chimaera* <i>Apristurus</i> sp. deepwater catshark* <i>Bathyraja</i> sp. skate Unidentified notacanthid spiny eel Unidentified halosaur <i>Serrivomer beanii</i> sawpalate eel* <i>Diastobranthus capensis</i> cuthroat eel* Unidentified cuthroat eel Unidentified alepocephalid smoothhead <i>Bathypterois dubius</i> spiderfish* <i>Coryphaenoides</i> sp. grenadier Unidentified morid codling <i>Anoplogaster cornuta</i> fangtooth* <i>Antigonia capros</i> deepbody boarfish* <i>Sebastes</i> sp. redfish <i>Zoarces atlanticus</i> eel-pout
ARTHROPODA	
<i>Neolithodes</i> sp. stone crab Unidentified spider crab assoc. w/ urchin	
ECHINODERMATA	
<i>Solaster</i> sp. sunstar <i>Porania</i> sp. sea star	

* identified by J. A. Moore, from commercial trawl 1997.

Results

Table 2 provides a list of the species collected during cruise DE00-11 and identified to date. We report a total of at least 274 species collected on or over Bear Seamount (Table 1 contains an additional 12 species not in Table 2 to give a total of at least 286 species known from Bear Seamount). The numbers of individuals for each species were recorded to give a qualitative sense of the relative abundances. For some very abundant animals (e.g. euphausiids, *Cyclothone*, *Sergestes* and *Pasiphaea*), only a small number of representative specimens were counted and kept, while for some fragmentary animals (e.g. hexactinellid sponges, *Lophelia pertusa* and *Lepidisis*) not all individuals were determinable. In addition, some ani-

mals were badly damaged by the trawl nets and difficult to identify (e.g. Myctophidae *indet.* and *Cyclothone* sp. *indet.*).

Preliminary identifications indicate that 115 fish species were captured (Table 2). Several new fish records for the area or rare species were collected (Moore *et al.*, 2003, including *Acromycter perturbator* (Congridae), *Diastobranthus capensis* (Synbranchidae), *Alepocephalus bairdii* (Alepocephalidae), *Mirognathus normani* (Alepocephalidae), *Bathygadus favosus* (Bathygadidae), *Nezumia longe-barbata* (Macrouridae), *Gaidropsarus argentatus* (Phycidae), *Dibranchius tremendus* (Ogcocephalidae), and *Kali indica* (Chiasmodontidae). The cephalopods comprised 26 species from 15 families, including one new

TABLE 2. Taxa collected on or over Bear Seamount during NMFS cruise DE00-11 from 2–7 Dec 2000.

PROTISTA		ANNELIDA (Continued)	
Agglutinated foraminiferans (3+ spp.) ¹	>20 ²	Serpulid with large triangular tubes	>4
		Serpulid with small round tubes	>40
PORIFERA		MOLLUSCA	
Boring sponge (in gorgonian base)	1	Nudibranch	1
Demospongae fragments	>6	<i>Calliotropis bairdii</i>	5
Small stoloniferous sponge	>4	<i>Pleurotomella</i> sp.	1
<i>Geodia</i> sp.	1	Gastropod sp. 1	1
Hexactinellid fragments	>6	Gastropod sp. 2	1
		Gastropod sp. 3	11
		Bivalve sp. 1	1
		Heteropods	>10
		Pteropods	>20
		<i>Clione</i> sp.	1
CNIDARIA		CEPHALOPODS	
Scyphozoans (8+ spp.)	>89	<i>Lampadioteuthis megaleia</i>	1
<i>Poralia rufescens</i>	1	<i>Abralia veranyi</i>	1
Siphonophore pieces	>8	<i>Abraliopsis hoylei</i>	5
<i>Marrus orthocanna</i>	1	<i>Pyroteuthis margaritifera</i>	1
Knobby pink anemone	2	<i>Pterygioteuthis gemmata</i>	1
Small white anemone	3	<i>Pterygioteuthis giardi</i>	1
<i>Amphianthus</i> or <i>Stephanauge</i> sp.	2	<i>Taningia danae</i>	1
Thecate hydroid attached to <i>Lophelia</i>	1	<i>Octopoteuthis megaptera</i>	1
<i>Epizoanthus paguriphilus</i>	46	<i>Gonatus fabricii</i>	2
<i>Caryophyllia ambrosia</i>	10	<i>Histioteuthis</i> sp.	4
<i>Flabellum alabastrum</i>	7	<i>Histioteuthis dofleini</i>	1
<i>Vaughanella margareta</i>	1	<i>Histioteuthis reversa</i>	2
<i>Lophelia pertusa</i>	5	<i>Bathyteuthis abyssicola</i>	1
<i>Acanthogorgia angustiflora</i>	2	<i>Brachioteuthis</i> sp.	1
<i>Paragorgia</i> sp.	5	<i>Illex illecebrosus</i>	2
<i>Swiftia?</i> sp.	1	<i>Chiroteuthis</i> sp.	2
<i>Lepidisis</i> sp.	>5	<i>Chiroteuthis spoeli</i>	1
<i>Anthoptilum grandiflorum</i>	9	<i>Chiroteuthis veranii</i>	2
<i>Funiculina</i> sp.	1	<i>Mastigoteuthis</i> sp.	5
<i>Anthomastis agassizii</i>	4	<i>Mastigoteuthis agassizi</i>	8
		<i>Mastigoteuthis magna</i>	25
		Cranchiidae indet.	1
		<i>Taonius pavo</i>	2
		<i>Teuthowenia megalops</i>	5
		<i>Stauroteuthis syrtensis</i>	2
		<i>Eledonella pygmaea</i>	1
		<i>Macrotritopus defilippi</i>	2
		<i>Graneledone verrucosa verrucosa</i>	2
		<i>Vampyroteuthis infernalis</i>	2
CHAETOGNATHA			
Chaetognaths	>3		
NEMERTEANS			
<i>Nectonemertes mirabilis</i>	2		
<i>Dinonemertes</i> cf. <i>investigatoris</i>	1		
ANNELIDA			
Polychaete burrowing in wood	3		
Polychaete in tubes within <i>Lophelia</i>	2		

¹ In a few instances, taxonomic experts indicated the number of species they could distinguish, without necessarily giving the explicit species names and these numbers of species are indicated below in parentheses following the taxonomic group name.

² For some very abundant animals (euphausids, *Cyclothone*, and *Pasiphaea*), only a small number of representative specimens were counted and kept, and for some or fragmentary animals (e.g. *Ophiomusium lymani* and *Lepidisis*) not all individuals were determinable. These are noted with a > sign before the actual number counted.

TABLE 2. (Continued). Taxa collected on or over Bear Seamount during NMFS cruise DE00-11 from 2–7 Dec 2000.

ARTHROPODA		ECHINODERMATA (Continued)	
<i>Syscenus atlanticus</i>	27	<i>Ophiomusim lymani</i>	>141
Bopyrid isopod (in <i>Glyphocrangon</i>)	1	Ophiacanthidae sp.	1
Hyperiid amphipods indet.	>49	<i>Ophiacantha</i> sp.	4
<i>Hyperia galba</i>	2	<i>Amphiophiura saurura</i>	8
<i>Phronima</i> sp.	2	<i>Homophiura tessellata</i>	9
Unidentified large red amphipod 1		<i>Ophiocten gracilis</i>	7
<i>Colossendeis colossea</i>	22	<i>Ophiura ljungmani</i>	36
<i>Scalpellum</i> sp. 1		Simple-armed basket star	1
<i>Plesiopenaeus edwardsianus</i>	10	<i>Echinus affinis</i>	159
<i>Acanthephyra</i> sp. (4 spp.)	>78	<i>Echinus</i> sp.	3
<i>Glyphocrangon sculpta</i>	>10	<i>Zygothuria lactea</i>	4
<i>Notostomus</i> sp.	4	<i>Phorosoma placenta</i>	20
<i>Pasiphaea</i> sp. (3 spp.)	>51	<i>Hygrosoma petersi</i>	9
<i>Gennadas</i> sp. (3–4 spp.)	>41		
<i>Sergestes</i> sp. (10 spp.)	>130	UROCHORDATA	
Homolids	2	Stalked tunicates	9
<i>Oplophorus</i> sp.	1	Thaliacean salps	>9
<i>Systellaspis</i> sp.	2	<i>Pyrosoma</i> sp.	>3
Euphausiids	>53		
<i>Pandalus carinatus</i>	1	PISCES	
<i>Stylopandalas</i> sp.	7	<i>Hydrolagus affinis</i>	4
<i>Polycheles granulatus</i>	12	<i>Apristurus manis</i>	14
<i>Steromastis nana</i>	6	<i>Apristurus profundorum</i>	5
Polychelid spiny form	2	<i>Etmopterus princeps</i>	2
<i>Munidopsis</i> sp.	8	<i>Raja bigelowi</i>	2
<i>Galathea rostrata</i>	1	<i>Aldrovandia</i> sp. indet.	13
<i>Munidopsis curvirostra</i>	1	<i>Aldrovandia affinis</i>	48
Mysids	>69	<i>Aldrovandia oleosa</i>	1
<i>Gnathophausia zoea</i>	>24	<i>Aldrovandia phalacra</i>	171
<i>Parapagurus pilosimanus</i>	46	<i>Halosauropsis macrochir</i>	4
<i>Neolithodes grimaldii</i>	1	<i>Notacanthus chemnitzii</i>	4
Copepod parasite on <i>Synbranchus</i> eel	1	<i>Polyacanthonotus challengeri</i>	1
		<i>Polyacanthonotus rissoanus</i>	3
ECHINODERMATA		<i>Ilyophis brunneus</i>	2
Crinoid fragments	2	<i>Diastobranchus capensis</i>	1
Elasipodid holothurian	6	<i>Synaphobranchus</i> sp.	17
<i>Psychropotes depressa</i>	1	<i>Synaphobranchus kaupii</i>	90
<i>Plutonaster agassizi</i>	6	<i>Derichthys serpentinus</i>	1
<i>Psilaster andromeda florae</i>	1	<i>Nessorhamphus ingolfianus</i>	2
<i>Cheiraster sepius</i>	6	<i>Nemichthys scolopaceus</i>	10
<i>Persephonaster</i> sp.	2	Congridae larvae	7
<i>Ceramaster granularis</i>	1	<i>Acromycter perturbator</i>	1
<i>Mediaster bairdi</i>	15	<i>Ariosoma</i> sp. Larvae	16
<i>Pseudarchaster parelii</i>	2	<i>Venefica procera</i>	2
<i>Neomorphaster forcipatus</i>	95	<i>Serrivomer beanii</i>	27
<i>Chondraster grandis</i>	3	<i>Eurypharynx pelacanoides</i>	11
<i>Henricia</i> sp.	1	<i>Bathylagus berycoides</i>	1
<i>Pteraster</i> sp.	1	<i>Bathylagus euryops</i>	12
<i>Solaster benedicti</i>	1	<i>Dolichopteryx</i> sp.	2
<i>Zoroaster</i> sp.	1	<i>Alepocephalus agassizi</i>	8
<i>Brisingia costata</i>	1	<i>Alepocephalus australis</i>	13
<i>Freyella microspina</i>	2	<i>Alepocephalus bairdii</i>	1
<i>Freyella elegans</i>	1		
<i>Asteroschema</i> sp.	2		

TABLE 2. (Continued). Taxa collected on or over Bear Seamount during NMFS cruise DE00-11 from 2–7 Dec 2000.

PISCES (continued)		PISCES (continued)	
<i>Alepocephalus</i> cf. <i>umbriceps</i>	1	<i>Lobianchia gemellari</i>	2
<i>Bajacalifornia megalops</i>	1	<i>Myctophum affine</i>	3
<i>Mirognathus normani</i>	1	<i>Nannobranchium atrum</i>	1
<i>Narctes stomias</i>	1	<i>Nannobranchium lineatum</i>	7
<i>Rouleina attrita</i>	1	<i>Nannobranchium mcdonaldi</i>	3
<i>Maulisia</i> sp.	2	<i>Nannobranchium lineatum</i>	7
<i>Cyclothone</i> sp. indet.	332	<i>Nannobranchium mcdonaldi</i>	3
<i>Cyclothone pallida</i>	1	<i>Notoscopelus resplendens</i>	2
<i>Cyclothone microdon</i>	>308	<i>Symbolophorus veranyi</i>	1
<i>Gonostoma bathyphilum</i>	32	<i>Dicrolene intronigra</i>	2
<i>Gonostoma elongatum</i>	58	<i>Penopus microcephalus</i>	1
<i>Chauliodus sloani</i>	34	<i>Porogadus miles</i>	4
<i>Argyropelecus aculeatus</i>	16	<i>Bassogigas gilli</i>	1
<i>Argyropelecus hemigymnus</i>	4	<i>Bathygadus favosus</i>	5
<i>Sternoptyx diaphana</i>	18	<i>Coryphaenoides alateralis</i>	2
<i>Sternoptyx pseudobscura</i>	1	<i>Coryphaenoides carapinus</i>	36
<i>Valenciennellus tripunctulatus</i>	1	<i>Coryphaenoides rupestris</i>	89
<i>Vinciguerrria nimbaria</i>	1	<i>Macrourus berglax</i>	25
<i>Malacosteus niger</i>	7	<i>Nezumia longebarbata</i>	5
<i>Aristostomias tittmani</i>	1	<i>Nezumia sclerorhynchus</i>	1
<i>Bathophilus pawneeii</i>	1	<i>Nezumia suilla</i>	2
<i>Borostomias antarcticus</i>	1	<i>Sphagemacrurus grenadae</i>	4
<i>Leptostomias longibarba</i>	1	<i>Trachonurus sulcatus</i>	2
<i>Melanostomias bartonbeani</i>	2	<i>Antimora rostrata</i>	20
<i>Photostomias gueneri</i>	4	<i>Gaidropsarus ensis</i>	2
<i>Stomias boa ferox</i>	20	<i>Gaidropsarus argenteus</i>	1
<i>Stomias affinis</i>	1	<i>Halargyreus johnsoni</i>	15
<i>Bathypterois phenax</i>	28	<i>Dibranchius tremendus</i>	1
<i>Bathypterois quadrifilis</i>	16	<i>Cryptopsaras couesii</i>	2
<i>Scopelosaurus lepidus</i>	1	<i>Melamphaes microps</i>	1
<i>Bathysaurus ferox</i>	3	<i>Melamphaes suborbitalis</i>	4
<i>Arctozenus rissoi</i>	1	<i>Poromitra megalops</i>	3
<i>Paralepis</i> sp.	2	<i>Scopelogadus beanie</i>	23
<i>Paralepis coregonoides</i>	1	<i>Scopeloberyx opisthopterus</i>	33
Myctophidae indet.	97	<i>Anoplogaster cornuta</i>	1
<i>Benthoosema glaciale</i>	73	<i>Cottunculus thomsoni</i>	1
<i>Ceratoscopelus maderensis</i>	18	<i>Howella brodei</i>	1
<i>Ceratoscopelus warmingii</i>	5	<i>Anthias nicholsi</i> larva	1
<i>Diaphus brachycephalus</i>	1	<i>Platyberyx opalescens</i>	1
<i>Diaphus dumerili</i>	9	<i>Lycodes atlanticus</i>	6
<i>Diaphus rafinesquii</i>	4	<i>Melanostigma atlanticum</i>	1
<i>Hygophum taaningi</i>	1	<i>Chiasmodon niger</i>	4
<i>Lampadena speculigera</i>	1	<i>Kali indica</i>	1
<i>Lampanyctus photonotus</i>	4	<i>Glyptocephalus cynoglossus</i> larvae	3
<i>Lobianchia dofleini</i>	30	<i>Poecilopsetta beani</i> larva	1

distributional record (i.e. *Chiroteuthis spoeli*) and several rarely-collected species. The crustacean species numbered at least 46 species and again included unusual species. For example, a redescription of a rare isopod species (*Systemus atlanticus*, previously known only from off Iceland) has been submitted for publi-

cation based in part on specimens from Bear Seamount (Kensley, 2003). Totals for other invertebrate species are pending laboratory identification, but number at least 87 species in 10 Phyla. Notable captures of other invertebrates include a solitary coral, *Vaughanella margaretata*, which represents the first record of this

species since its original description over one hundred years ago (S. Cairns, National Museum of Natural History, Smithsonian Institution, Washington, D.C., USA, pers. comm.). We have also been informed of a number of new distributional records and an undescribed species of gorgonian (F. Bayer, National Museum of Natural History, Smithsonian Institution, Washington, D.C., USA, pers. comm.)

Only two fish species of potential commercial importance were encountered: *Coryphaenoides rupestris* and *Macrourus berglax*. *Coryphaenoides rupestris*, the roundnose grenadier, was caught at 4 stations at depths of 1 100–1 800 m. They ranged from 47–103 cm total length, with an average length of 85 cm total length. A total of 89 individuals were caught, with the highest catch of 46 individuals at one station, weighing a total of 65.1 kg. *Macrourus berglax*, the roughhead grenadier, was captured at 5 stations, also at depths of 1 100–1 800 m. They ranged from 31–104 cm total length, with an average length of 50 cm. Twenty-five individuals were caught, with a highest single catch of 10 individuals.

The only commercially exploited cephalopod captured, *Illex illecebrosus*, consisted of just two individuals. Only a few exploited or potentially exploitable crustaceans were taken from Bear Seamount (a few isolated individuals of scarlet shrimp, *Plesio-peneaus edwardsianus*, several glass shrimps, *Pasiphaea* spp., and the porcupine stone crab, *Neolithodes grimaldii*; Pohle *et al.*, 1992).

Discussion

Our results have some bearing on statements made by Wilson and Kaufman (1987) in their review of seamount biota. For instance, they state that although ophiuroid brittlestars are widespread and common in deep-sea sedimentary environments, they have been reported from very few seamounts. Our survey collected a number of ophiuroid species and found the brittlestar *Ophiomusium lymani* to be common in all the benthic trawls. Archived photos of other seamounts in the NES chain also show numerous brittlestars on soft sediment, on rocks, and perched in the fronds of gorgonians (see Fig 9 in Houghton *et al.*, 1977).

Wilson and Kaufman (1987) also state that few of the reported seamount crustaceans are truly benthic. However, we collected a number of benthic crustacean species from Bear Seamount. A large spider crab, *Neolithodes grimaldii*, and numerous hermit crabs,

Parapagurus pilosimanus, were taken, as were a variety of polychelid and galatheid crustaceans. Also, a gooseneck barnacle (*Scalpellum* sp.) was collected attached to a small glacial pebble. Archived photos from the two *Alvin* dives on Bear Seamount show a *Neolithodes* in a shallow dish-like depression, an unidentified small spider crab associated with an urchin, and large red shrimp, possibly *Plesio-peneaus edwardsianus*, just inside the mouths of small burrows in the soft sediment.

With regards to seamount fish fauna, Wilson and Kaufman (1987) state that fishes are associated with seamounts elsewhere, but have not been reported from the western North Atlantic seamounts. This was true at the time of that paper. Since then, Vinnichenko (1997) published a list of the fishes found on the Corner Rise Seamounts (about 300 km east to the eastern end of the NES). That faunal list compares well with our list of fishes from Bear Seamount. The main difference is that the Corner Rise features many species more typically found in the deep waters of the eastern North Atlantic, while our list has many more western North Atlantic deepwater species. However, a few eastern Atlantic species (e.g. *Hydrolagus pallida* and *Bathypterois dubius*) were found on Bear Seamount and may indicate that the NES chain provides a westward route of dispersal for at least some eastern Atlantic species.

Large pieces of live *Lophelia pertusa* were collected from two of the bottom hauls indicating that deepwater coral banks may exist on this seamount. Evidence of *Lophelia* sp. has also been noted from other areas of NES (Pratt and Thomson, MS 1962; Heitzler *et al.*, MS 1977). This species has been associated with very diverse communities elsewhere in the North Atlantic (Jensen and Frederiksen, 1992; Mortensen *et al.*, 1995; Rogers, 1999). Similarly diverse communities may also be present on Bear Seamount. We can note the following invertebrates found in direct association with the *L. pertusa* trawled from Bear Seamount: a large polychaete worm living in tubular spaces within the coral colony, attached solitary coral, *Caryophila ambrosia*, small serpulid worm tubes, a thecate hydroid, and the octocoral *Swiftia*.

Even though Bear Seamount rises up out of the continental slope of Georges Bank, it apparently provides sufficiently unusual habitats or conditions that allow various species not generally reported from the nearby slope to live on the seamount. Three of us have

worked on a checklist of deepwater fish species for all of New England (Moore *et al.*, 2003). The trawls on Bear Seamount resulted in six new additions to that checklist. Our regional list of deep-sea fishes recorded about 591 species in the vicinity of Georges Bank, from the slope to the abyssal plain, whereas we report 115 fish species from Bear Seamount. It is clear that this investigation of Bear Seamount is quite preliminary and that many more species of fishes and invertebrates will likely be found associated with the seamount. Because of the more unusual habitats on the seamount, it is also quite possible that more new additions to the region will also be found on the seamount.

Similarly, the cephalopods from this survey add to the slope fauna recently compiled by Vecchione (2000) and Vecchione and Pohle (2002). Photographs of some of these unusual cephalopods have been added to the cephalopod section of the Tree of Life project on the World Wide Web at:

<http://tolweb.org/tree/eukaryotes/animals/mollusca/cephalopoda/cephalopoda.html>

The numerous novel records of species associated with the seamount in our study could indicate that there is some degree of separation of the seamount fauna from the nearby continental slope, or our ignorance of what lives in the deep sea may still be so great that a single survey can add much new information, despite many deepwater investigations in the region over the last 130 years.

Acknowledgements

We are grateful to the following taxonomic experts who helped identify specimens from the various groups indicated: Susan Richardson (Foraminifera and Xenophyophora), Florida Atlantic University, Larry Madin (Scyphozoa) Woods Hole Oceanographic Institution, Steve Cairns (Scleractinia), Ted Bayer (Octocorallia), Ed Petuch (Bivalvia and Gastropoda), Jon Norenburg (Nemertea), Brian Kensley (Arthropoda), Dave Pawson (Echinodermata) all from Smithsonian Institution, Eric Lazo-Wasem (various invertebrates) Peabody Museum, Yale University, and Tomio Iwamoto (Macrouridae) California Academy of Science. Our thanks go to the captain and crew of the R/V *Delaware*, who worked hard, despite adverse seas and equipment failures, to make this cruise so successful. Thanks also go to Capt. Walter Allyn of the F/V *Matthew Melissa* for providing specimens from the exploratory trawling in 1997.

References

- AUSTER, P. J., R. J. MALTESTA, R. W. LANGTON, L. WATLING, P. C. VALENTINE, C. L. S. DONALDSON, E. W. LANGTON, A. N. SHEPARD, and I. G. BABB. 1996. The impacts of mobile fishing gear on seafloor habitats in the Gulf of Maine (Northwest Atlantic): Implications for conservation of fish populations. *Rev. Fish. Sci.*, **4**(2): 185–202.
- DAYTON, P. K., S. F. THRUSH, M. T. AGARDY, and R. J. HOFMAN. 1995. Environmental effects of marine fishing. *Aquat. Conserv.: Mar. Freshw. Ecosyst.*, **5**: 205–232.
- HAMILTON, P., S. MCDOWELL, E. WADDELL, D. REDFORD, and D. PABST. 1996. Slope sea currents during the 1990–1991 106-mile site investigations. *J. Marine Env. Eng.*, **2**: 203–225.
- HEIRTZLER, J. R., P. T. TAYLOR, R. D. BALLARD, and R. L. HOUGHTON. MS 1977. The 1974 Alvin dives on Corner Rise and New England Seamounts. *WHOI Rep.*, WHOI-77-08, 53 p.
- HEIRTZLER, J. R., P. T. TAYLOR, R. D. BALLARD, and R. L. HOUGHTON. 1977. A visit to the New England Seamounts. *Amer. Sci.*, **65**: 466–472.
- HOGG, N. G., R. S. PICKART, R. M. HENDRY, and W. J. SMETHIE, JR. 1986. The Northern Recirculation Gyre of the Gulf Stream. *Deep-Sea Res.*, **33**(9A): 1139–1165.
- HOUGHTON, R. L., J. R. HEIRTZLER, R. D. BALLARD, and P. T. TAYLOR. 1977. Submersible observations of the New England Seamounts. *Naturwissenschaften*, **64**: 348–355.
- JENSEN, A. and R. FREDERIKSEN. 1992. The fauna associated with the bank-forming deepwater coral *Lophelia pertusa* (Scleractinaria) on the Faroe Shelf. *Sarsia*, **77**: 53–69.
- KENSLEY, B. 2003. Redescription and distribution of two species of *Syscenus* (Crustacea, Isopoda) in the North Atlantic. *Sarsia*, **88** (in press).
- KOSLOW, J. A. 1997. Seamounts and the ecology of deep-sea fisheries. *Amer. Sci.*, **85**: 168–176.
- KOSLOW, J. A. and K. GOWLETT-HOLMES. MS 1998. The seamount fauna off southern Tasmania: benthic communities, their conservation and impacts of trawling. Final report to Environment Australia and the Fisheries Research and Development Corporation. CSIRO, Hobart, Tasmania (Australia) 104 p.
- KOSLOW, J. A., G. W. BOEHLERT, J. D. M. GORDON, R. L. HAEDRICH, P. LORANCE, and N. PARIN. 2000. Continental slope and deep-sea fisheries: implications for a fragile ecosystem. *ICES J. Mar. Sci.*, **57**: 548–557.
- MOORE, J. A. and P. M. MACE. 1999. Challenges and prospects for deep-sea finfish fisheries. *Fisheries*, **24**(7): 22–23.
- MOORE, J. A., K. E. HARTEL, J. E. CRADDOCK, and J. K. GALBRAITH. 2003. An annotated checklist of deepwater fishes from off New England, with 110 range extensions and 81 new records off New England. *North-east Nat.*, **10**(2): 159–248.
- MORTENSEN, P. B., M. HOVLAND, T. BRATTEGARD,

- and R. FARESTVEIT. 1995. Deep water bioherms of the scleractinian coral *Lophelia pertusa* (L.) at 64°N on the Norwegian shelf: structure and associated megafauna. *Sarsia*, **80**: 145–158.
- POHLE, G., T. J. KENCHINGTON, and R. G. HALLIDAY. 1992. Potentially exploitable deepwater resources off Atlantic Canada. *Can. Tech. Rep. Fish. Aquat. Sci.*, **1843**: 1–79.
- PRATT, R. M. and S. L. THOMPSON. MS 1962. Report on Atlantis cruises #280–281, June–July 1962. *WHOI Rep.*, WHOI-62-40, 48 p.
- PROBERT, P. K., D. G. MCKNIGHT, and S. L. GROVE. 1997. Benthic invertebrate bycatch from a deep-water trawl fishery, Chatham Rise, New Zealand. *Aquat. Conserv.: Mar. Freshw. Ecosyst.*, **7**: 27–40.
- ROGERS, A. D. 1994. The biology of seamounts. *Adv. Mar. Biol.*, **30**: 305–350.
- ROGERS, A. D. 1999. The biology of *Lophelia pertusa* (Linnaeus, 1758) and other deep-water reef-forming corals and impacts from human activity. *Int. Rev. Hydrobiol.*, **84**(4): 315–406.
- SLEEP, N. H. 1990. Montereian hotspot track: a long-lived mantle plume. *J. Geophys. Res.*, **95**(B13): 21,983–21,990.
- UCHUPI, E, J. D. PHILLIPS, and K. E. PRADA. 1970. Origin and structure of the New England Chain. *Deep-Sea Res.*, **17**: 483–494.
- VASTANO, A. C. and B. A. WARREN. 1976. Perturbations of the Gulf Stream by Atlantis II seamount. *Deep-Sea Res.*, **23**: 681–694.
- VECCHIONE, M. 2000. Cephalopods of the continental slope east of the United States. in: G. Sedberry (ed.) Islands in the Stream. Oceanography and Fisheries of the Charleston Bump. *Amer. Fish. Soc. Sym.*, **25**: 153–160.
- VECCHIONE, M. and G. POHLE. 2002. Midwater cephalopods in the western North Atlantic Ocean of Nova Scotia. *Bull. Mar. Sci.*, **71**: 883–892.
- VINNICHENKO, V. I. 1997. Russian investigations and deep water fishery on the Corner Rising Seamount in Subarea 6. *NAFO Sci. Council Studies.*, **30**: 41–49.
- WILSON, R. R., JR. and R. S. KAUFMAN. 1987. Seamount biota and biogeography. In: Seamounts, islands, and atolls. B. H. Keating, P. Fryer, R. Batiza, and G. W. Boehlert (eds.). *Geophys. Monogr.*, No. 43, p. 355–377.
- ZEIGLER, J. M. MS 1955. Seamounts near the eastern coast of North America. *WHOI Rep.*, WHOI-55-17, 36 p.
-